

A CATALOGUE OF PHILIPPINE HYMENOPTERA, WITH DESCRIPTIONS OF NEW SPECIES.

By ROBERT E. BROWN, S. J., *Manila Observatory*.

In 1885, Don Ramon Jordana published in Madrid a work entitled "Bosquejo geográfico é histórico-natural del Archipiélago Filipino," in which he tabulated all the species then known to belong to the Philippine fauna. The order *Hymenoptera* was very poorly represented, no more than some thirty species being given as having been met with in the Philippines and none of these were characteristic. Ten years later, on the occasion of the exposition which was held in Manila in 1895, Father Castro de Elera, O. P., published another catalogue of the fauna of the Islands, which he entitled "Catálogo sistemático de toda la fauna de Filipinas conocida hasta el presente." This work is much more pretentious and complete than that of Señor Jordana, though it is somewhat marred by the inclusion of many exotic forms which were contained in the Museum of the University of Saint Thomas; and although Father Elera explains this in the subtitle and notes the exotic species in the body of the work, still, unless one has this constantly before one's mind, the impression is gained that the number of Philippine species is greater than it really is. In the catalogue of the *Hymenoptera* which is given in the second volume, pages 238-247, Father Elera enumerates fifty-four species as representing the known Philippine insects of this order. This is not quite correct, as he had inadvertently overlooked a paper published by Professor Emery in the Annals of the Entomological Society of France in 1893 in which forty-one species of ants are recorded from the Island of Luzon, principally from Manila and Antipolo.

This then brought the *Hymenoptera* known in the Archipelago up to the year 1895 to the number of ninety-five.

In 1903, Rev. William Stanton, S. J., at that time Assistant Director of the Philippine Weather Bureau, began to study in his spare time the *Hymenoptera* which he collected in the garden attached to the Observatory. He discovered many forms which he did not recognize, and, being unable to classify them owing to the lack of literature on the subject, he sent his specimens to Dr. William Ashmead, of the National Museum, Washington, who found that the great majority of the species were entirely new to science and that almost all the remainder were not before recorded from the Islands.

Father Stanton continued the work till April, 1904, when he was obliged to return to the United States, but in the short time he had been collecting he had discovered no less than five new genera and sixty-eight new species.

The descriptions of the earlier of these new species were published by Dr. Ashmead in the Journal of the New York Entomological Society and of the later collections in the Canadian Entomologist and the Proceedings of the United States National Museum.

When Father Stanton left the Philippines, the writer continued his work and as a consequence the number of new genera and new species has greatly increased; in fact, from December to June, which constitutes the best collecting season, the average of new species was one per day and of new genera three every two months.

The number of species in the adjoined catalogue is 458. Many are included which were not contained in Dr. Ashmead's check list, as he had apparently overlooked several recorded by Kohl, Ritsema, and Radoszowski, while species not before recorded from the Philippines are being constantly discovered. However, this number does not represent the complete hymenopterous fauna, for the writer has in his possession some thirty-nine new species which have not as yet been described, while the Entomological section of the Bureau of Science has many more undescribed forms.

After the name of the species discovered since 1903 I have placed the name of the collector and in the case of those found by Father Stanton and the writer, M. O., to show that they were obtained from the garden of the Manila Observatory. At the end of the catalogue will be found a list of some of the hosts from which a few of the new species were bred in the Observatory.

Superfamily III. VESPOIDEA.

Family XXVII. CEROPALIDÆ.

SUBFAMILY III. APORINÆ.

Tribe I. ANOPLIINI.

TACHYPOMPILUS Ashmead.

Tachypompilus ashmeadii sp. nov. ♀ Length, 14–16 millimeters. Head, thorax, and part of abdomen black, last 3 segments of abdomen brick red, covered with sparse pubescence, the pubescence consisting of straight, longish hairs; head smooth, pronotum lightly rugose, scutellum, metanotum and abdomen finely punctate, shining. Clypeus truncate, smooth. Scape stout, first joint of flagellum twice as long as the second, front face long, with a deep medial furrow. Anterior face of metathorax convex, posterior face concave with the angles obtusely dentate. Wings hyaline with nervules fuscous, median and submedian cells of equal length. Three cubital cells, first almost triangular, second quadrangular and third very large; hind tibia sparsely armed with spines, first joint of hind tarsi as long again as any of the rest, claws reddish. Manila.

Family XXXI. CHRYSIDIDÆ.

Subfamily II. CHRYSIDINÆ.

CHRYYSIS Linn.

Chrysis miri sp. nov. ♂ Length, 6.5 millimeters. Near to *C. mendicalis* Cameron. Head, thorax, first and third abdominal segments, light metallic blue; space around ocelli, a few punctures on collar and pronotum, medial area of scutellum and second abdominal segment purple; round spot and apical margin of third segment purple, with a transverse line on basal area of second segment, dark purple.

Antennæ black, the scape in front and two basal joints of flagellum light bronze green; coxae and femora of front legs metallic green; tibiæ slightly green; tarsi dark brown, tibia with slight cinerous pubescence; hind legs green, tarsi brownish. Head and thorax with closely set, coarse punctures, with smaller ones scattered up and down, the punctures larger on posterior portion of thorax, smaller on the abdomen and very symmetrically arranged. The ventral segments very finely punctate, with a brown medial longitudinal ridge along all the segments.

Antennæ thick, the third joint of the flagellum as long again as the second. Pronotum broad, very much rounded anteriorly, medial area of mesonotum separated from the lateral areas by longitudinal rows of purple punctures; scutellum and post-scutellum convex, posterior portion of post-scutellum very much rounded.

First abdominal segment with a medial wide and deep depression anteriorly. Apical segment of third segment with 3 teeth, one on each angle and one in the middle, the intervals between the teeth sinuate.

Family XLII. MUTILLIDÆ.

Subfamily I. MUTILLINÆ.

MUTILLA Linn.

Mutilla manilensis sp. nov. This species comes very near to *M. vicinissima* Gribodo but is differentiated by the size of the maculæ and the bands of pubescence on the third and fifth segments. ♀ Length, 7.5 millimeters. Head broader than thorax. Head and thorax coarsely, abdomen finely, punctured. Scape, femora, head posteriorly, thorax and abdomen densely pubescent. Head, antennæ, abdomen and legs black, thorax red; pubescence on abdomen cinerous, on legs and antennæ whitish.

The second abdominal segment above with 2 small, round, silvery maculæ; the third and fifth segments covered with transverse bands of silvery pubescence. The first ventral segment with a prominent longitudinal carina, the fourth and fifth segments coarsely punctate posteriorly and striated anteriorly.

Mutilla parva sp. nov. Near to *M. decora* Smith. ♀ Length, 4.5 millimeters. Head and abdomen finely, and thorax coarsely, punctate and sparsely pubescent, the pubescence consisting of thin, long, curved hairs. Eyes, large, oval; thorax very much rounded, abrupt anteriorly and with a steep slope posteriorly; head and antennæ black, except the scape which is rufous covered with whitish pubescence; legs, except the femora, and abdomen black, thorax red.

A small oval silvery-white spot in the middle above on the second abdominal segment, a transverse band of silvery-white pubescence which is interrupted medially on its posterior margin, and the third segment also covered with the white pubescence. First, third, and fourth ventral segments strongly longitudinally carinate.

CATALOGUE OF PHILIPPINE HYMENOPTERA.

Order HYMENOPTERA.

Suborder I. **Heterophaga** Ashmead.Superfamily I. **APOIDEA**.Family I. **APIDÆ**.*Megapis dorsata* Fabricius. *zonata* Smith.*Apis indica* Fabricius. *mellifera* Linnaeus. *nigrocineta* Smith. *unicolor* Latreille.*Mierapis florea* Fabricius.*Trigone biroi* Friese.*Melipona laeviceps* Smith.Family II. **BOMBIDÆ**.*Bombus mearnsi* Ashmead; E. A. Mearns.Family V. **ANTHOPHORIDÆ**.*Anthophora cingulata* Fabricius. *zonata* Linnaeus.Family VI. **NOMADIDÆ**.*Crocisa emarginata* Lepeletier. *lamprosoma* Boisduval. *nitidula* Fabricius.*Nomada lusea* Smith. *philippinensis* Vachal.*Nomia dimidiata* Vachal.Family VII. **CERATINIDÆ**.*Ceratina compacta* Smith. *hieroglyphica* Smith. *imperialis* Ashmead; M. O. *philippinensis* Ashmead; M. O. *sexmaculata* Smith. *tropica* Ashmead; M. O.*Allodape jueunda* Smith. *minutior* Ashmead; M. O. *philippinensis* Ashmead; M. O.Family VIII. **XYLOCOPIDÆ**.*Xylocopa aestuans* Linnaeus. *adusta* Perez. *amauroptera* Perez. *bombiformis* Smith. *bryorum* Fabricius.*Xylocopa dissimilis* Lepeletier. *euchlora* Perez. *fuliginosa* Perez. *ghilianii* Gribodo. *latipes* Fabricius. *occipitalis* Perez. *philippinensis* Smith. *sonorina* Smith. *sulcifrons* Perez. *rufescens* Smith. *tricolor* Ritsema. *trifasciata* Gribodo. *vachali* Perez.*Platynopoda latipes* Drury. *tenuicornis* Westwood.Family IX. **MEGACHILIDÆ**.*Megachile atrata* Smith. *clotho* Smith. *hera* Bingham. *lachesis* Smith. *laticeps* Smith. *robbii* Ashmead; M. L. Robb.*Ctenoplectra vagans* Cockerell.Family X. **STELLIDÆ**.*Coelioxys manilae* Ashmead; M. O. *philippinensis* Bingham.Family XII. **ANDRENIDÆ**.*Hoplonomia quadrifasciata* Ashmead; M. O.*Paranomia stantoni* Ashmead; M. O.*Halictus manilae* Ashmead; M. O. *robbii* Ashmead; M. L. Robb.Family XIV. **PROSOPIDÆ**.*Prosopis tagala* Ashmead; C. S. Banks.Superfamily II. **SPHECOIDEA**.Family XV. **OXYBELIDÆ**.*Notoglossa banksi* Ashmead; C. S. Banks.

Family XVI. CRABRONIDÆ.

Subfamily III. CRABRONINÆ.

Crabro quadriceps Bingham.
venans Kohl.

Subfamily IV. RHOPALINÆ.

Dasyproctus philippensis Ashmead;
M. O.

Rhopalum albocollaris Ashmead; M. O.

Family XVII. PEMPHREDO-NIDÆ.

Psen algii Ashmead; M. O.

Family XVIII. BEMBICIDÆ.

Bembex borei Handlsch.
melancolica Smith.

Family XIX. LARRIDÆ.

Subfamily I. LARRINÆ.

Notogonia laboriosa Smith.
manilae Ashmead; M. O.
subtessalata Smith.

Larra aurata Fabricius.

Threosphex stantoni Ashmead; M. O.

Subfamily IV. PISONINÆ.

Pison lagunae Ashmead; M. O.; P. L. Stangl.
punctatus Ashmead; M. O.
Pisonitus argenteus Ashmead; P. L. Stangl.
Pisonoides browni Ashmead; M. O.

Family XX. PHILANTHIDÆ.

Subfamily I. CERCERINÆ.

Cerceris vafra Bingham.

Family XXI. TRYPOXYLIDÆ.

Trypoxylon bicolor Smith.
elongatum Ashmead; M. O.
philippensis Ashmead; M. O.

Family XXII. MELLINIDÆ.

Megalomma quadricinctum Ashmead;
M. O.

Family XXV. SPHECIDÆ.

Subfamily I. SPHECINÆ.

Sphex argentata Dahlbom.
aurulentus Fabricius.
(a) *var. ferrugineus* Lepeletier.
(b) *var. lineolus* Lepeletier.
sericeus Fabricius.
sulciscuta Gribodo.
splendidus Kohl.
umbrosus Christ.
(a) *var. plumbiferus* Costa.
(b) *var. rufipennis* Fabricius.

Chlorion lobatum Fabricius.

Subfamily II. AMMOPHILINÆ.

Ammophila atripes Smith.
coronata Costa.
superciliaris Saussure.

Subfamily III. SCELIPHRONINÆ.

Sceliphron bengalensis Dahlbom.
madraspatanum Fabricius.
var. conspicillatum Costa.
violaceum Fabricius.

Chalybion violaceum Dahlbom.

Family AMPULICIDÆ.

Subfamily AMPULICINÆ.

Ampulex compressa Fabricius.
laevigata Kohl.
sonneratii Kohl.

Superfamily III. VESPOIDEA.

Family XXVII. CEROPALIDÆ.

Subfamily I. PEPSINÆ.

Salius flavus Fabricius.
graphicus Smith.
pulchripennis Smith.
Hemipepsis tagala Gribodo.
Pallosoma fulgidipennis Saussure.
Pseudosalius bipartitus Lepeletier.
Caligurgus sericosoma Smith.

Subfamily II. AGENIINÆ.

Macromeris argentifrons Smith.
splendida Lepeletier.
violacea Lepeletier.

Pseudagenia femorata Ashmead; M. O.
imitator Ashmead; M. O.
rufofemorata Ashmead;
 M. O.

unifasciata Ashmead; M. O.

Spilopompilus stantoni Ashmead; M. O.

Agenia cingulata Ashmead; M. O.

Subfamily III. APORINÆ
(Pompilinæ).

Tribe I. *Anopliini.*

Tachypompilus ashmeadii Brown; M. O.

Family XXVIII. VESPIDÆ.

Subfamily I. VESPINÆ.

Vespa analis Fabricius.

annulata Smith.

cineta Fabricius.

deusta Lepeletier.

luctuosa Saussure.

nigripennis Saussure.

philippensis Saussure.

Provespa doryloidea Saussure.

Subfamily II. POLISTINÆ.

Polistella manilensis Saussure.

Polistes dubius Saussure.

hebraeus Fabricius.

philippensis Saussure.

Icaria cagayanensis Ashmead; M. O.

philippensis Saussure.

speciosa Saussure.

Family XXIX. EUMENIDÆ.

Subfamily IV. EUMENINÆ.

Eumenes conica Fabricius.

curvata Saussure.

fulvipennis Smith.

gracilis Saussure.

pomiformis Fabricius.

Rhynchium argentatum Fabricius.

atrum Saussure.

Lionotus dyscherus Saussure.

punctum Saussure.

xanthozonatus Ashmead; M. O.

Ancistrocerus bizonatus Boisduval.

Odynerus punctum Saussure.

Family XXXI. CHRYSISIDÆ.

Subfamily II. CHRYSISIDÆ.

Stilbum cyanurum Fabricius.

amethystinum Fabricius.

splendidum Fabricius.

Chrysis fuscipennis Brullé.

miri Brown; M. O.

oculata Fabricius.

var. sumptuosa Gribodo.

triacantha Moscary.

Trichrysis aspera Brullé.

Hedychrum stantoni Ashmead; M. O.

Family XXXII. BETHYLIIDÆ.

Subfamily I. BETHYLINÆ.

Dissomphalis tibialis Ashmead; M. O.

Epyris tagala Ashmead; C. S. Banks.

Giniozus philippensis Ashmead; M. O.

Subfamily II. DRYININÆ.

Dryinus brownii Ashmead; M. O.

stantoni Ashmead; M. O.

Family XXXIII. TRYGONA-
 LIDÆ.

Trigonalyx lachrymosa Westwood.

Family XXXVI. SCOLIIDÆ.

Discolia aureipennis Lepeletier.

erratica Smith.

modesta Smith.

Scolia aurulenta Smith.

capitata Guerin.

grossa Fabricius.

manilae Ashmead; M. O.

modesta Smith.

procera Illiger.

var. bimaculata Gribodo.

pseudoforaminata Gribodo.

quadrifasciata Fabricius.

quadripustulata Fabricius.

rubiginosa Fabricius.

scutellaris Gribodo.

whiteheadii Bingham.

Subfamily II. ELIDINÆ.

- Elis albicollis* Christ.
annulata Fabricius.
aureicollis Lepeletier.
aurulenta Smith.
grossa Smith.
luctuosa Smith.
quadrifasciata Fabricius.
reticulata Cameron.
Liacos analis Fabricius.

Family XXXVII. TIPHIIDÆ.

- Tiphia compressa* Smith.
intrudens Smith.

Family XLII. MUTILLIDÆ.

Subfamily II. MUTILLINÆ.

- Mutilla accedens* Sichel.
analis Lepeletier.
dimidiata Lepeletier.
discreta Cameron.
luzonica Radoszhowski.
maculosofasciata Saussure.
manilensis Brown; M. O.
parva Brown; M. O.
philippinensis Smith.
semperi Ashmead; M. O.
suspiciosa Smith.
Trogaspidia bicolor Ashmead; C. S. Banks.
minor Ashmead; C. S. Banks.

Superfamily IV. FORMICOIDEA.

Family XLIV. PONERIDÆ.

Subfamily I. PONERINÆ.

- Ponera luteipes* Mayr.
Microponera nigriiceps Ashmead; M. O.

Subfamily II. PACHYCONDYLINÆ.

- Diacamma geometricum*.
var. viridipurpureum Emery.
versicolor Smith.
Odontoponera denticulata Smith.
transversa Smith.
Bothroponera glabripes Emery.

Family XLV. ODONTOMACHUS.

- Odontomachus haematodes* Linnaeus.
infandus Smith.
papuans Emery.

Family XLVI. MYRMICIDÆ.

- Subfamily I. PSEUDOMYRMICINÆ.
Sima allaborans Walker.

Subfamily II. MYRMICINÆ.

- Monomorium destructor* Jerdon.
Pheidole simoni Emery.
Cremastogaster bicolor Mayr.
crassicornis Emery.
longiclava Emery.
ochracea Mayr.
semperi Emery.
simoni Emery.
Pheidologeton diversus Jerdon.
pygmaeus Emery.
var. albipes Emery.
Plagiolepis longipes Jerdon.
Tetramorium guinensis Talrimis.
pacificum Mayr.
subsp. subscabrum Emery.
Solenopsis geminata Fabricius.

Family XLVIII. DOLICODERIDÆ.

- Dolichoderus bituberculatus* Mayr.
tuberculatus Mayr.
Technomyrmex albipes Smith.
Tapinoma melanocephalum Fabricius.

Family XLIX. FORMICIDÆ.

Subfamily II. CAMPONOTINÆ.

- Camponotus cinerascens* Fabricius.
gigas Latreille.
nigricans Roger.
pallidus Smith.
platypus Roger.
pennsylvanicus De Geer *var. japonicus* Mayr.
quadrisectus Smith.
Oecophylla smaragdina Fabricius.
var. subnitida Emery.
Calobopsis albocincta Ashmead; M. O.
pubescens Mayr.
Dinomyrmex gigas Latreille.

Polyrhachis abdominalis Smith.
aciulata Smith.
argentia Mayr.
armata Le Guillou.
bicolor Smith.
bihamata Drury.
cyaniventrus Drury.
dives Smith.
maligna Smith.
mayri Roger.
philippinensis Smith.
pubescens Mayr.
rastellata Smith.
sexspinosa Latreille.
trinax Roger.
subsp. javana Mayr.
subsp. sayonensis Forel.

Subfamily III. FORMICINÆ.

Plagiolepis longipes Jerdon.
Aphomyrmex emeryi Ashmead; M. O.
Formica quadrisecta Smith.
rubra Fabricius.

Superfamily V. PROCTOTRYPOIDEA.

Family LIV. DIAPRIIDÆ.

Diapria manilae Ashmead; M. O.
palida Ashmead; M. O.
philippinensis Ashmead; M. O.
tagala Ashmead; M. O.
Galesus luzonicus Ashmead; M. O.

Family LV. CERAPHRONIDÆ.

Ceraphron manilae Ashmead; M. O.

Family LVI. SCELIONIDÆ.

Subfamily III. TELEASINÆ.

Prósacantha roberti Ashmead; M. O.

Subfamily IV. SCELIONINÆ.

Macroteleia manilensis Ashmead; M. O.
Opithacanta manilae Ashmead; M. O.
nigrilavata Ashmead; M. O.
Telenomus catacanthae Ashmead; M. O.
Hoploteleia pacifica Ashmead; M. O.
Hadronotus flavipes Ashmead; M. O.
luteipes Ashmead; M. O.
Seilio philippinensis Ashmead; M. O.

Family LVII. PLATYGASTERIDÆ.

Anopedias luzonicus Ashmead; M. O.

Superfamily VI. CYNIPOIDEA.

Family LVIII. FIGITIDÆ.

Subfamily V. EUCOELINÆ.

Pentamerocera pacifica Ashmead; M. O.
Hexamerocera kiefferi Ashmead; M. O.
philippinensis Ashmead; M. O.
Kleidotoma philippinensis Ashmead; M. O.

Subfamily VI. XYSTINÆ.

Loboscelidia rufescens Westwood.

Superfamily VII. CHALCIDOIDEA.

Family AGAONIDÆ.

Subfamily I. AGAONINÆ.

Kradibia brownii Ashmead; M. O.

Family LXI. TORYMIDÆ.

Subfamily I. IDARNINÆ.

Sycoryctes philippinensis Ashmead; M. O.

Philotrypesis fieicola Ashmead; M. O.

Subfamily IV. PODAGRIONINÆ.

Podagrion philippinensis Ashmead; M. O.

Subfamily V. MEGASTIGMINÆ.

Megastigmus immaculatus Ashmead; M. O.

Family LXII. CHALCIDIDÆ.

Subfamily I. LEUCOSPIDINÆ.

Leucospis regalis Westwood.

Subfamily II. CHALCIDINÆ.

- Chaleis* *albotibialis* Ashmead; M. O.
argentifrons Ashmead; M. O.
banksi Ashmead; C. S. Banks.
obseurata Walker.
prodeniae Ashmead; M. O.
pulchripes Holmgren.
xerxena Walker.
Stomatocera *sulcata* Ashmead; M. O.
Arretocera *stantoni* Ashmead; M. O.
Neochaleis *tarsalis* Walker.
Haltichella *ludlowae* Ashmead; Miss
 C. S. Ludlow.
nasuta Holmgren.
validicornis Holmgren.
Dirhinus *anthracia* Walker.
auratus Ashmead; M. O.

Family LXIII. EURYTOMIDÆ.

- Eurytoma* *albotibialis* Ashmead; C. S.
 Banks.
banksi Ashmead; C. S.
 Banks.
carinatifrons Ashmead; M. O.
manilae Ashmead; M. O.
systolodes Ashmead; M. O.

Family LXV. EUCHARIDÆ.

- Chalcura* *aegineta* Walker.
nasua Walker.

Family LXVI. MISCOGASTE-
RIDÆ.

Subfamily III. MISCOGASTERINÆ.

- Omerocerus* *pallidipes* Ashmead; M. O.
Nesotoxeuma *petiolata* Ashmead; M. O.

Family LXVII. CLEONYMIDÆ.

Subfamily II. CLEONYMINÆ.

- Epistenia* *ania* Walker.
feretius Walker.

Family LXVIII. ENCYRTIDÆ.

Subfamily I. EUPELMINÆ.

- Metapelma* *gloriosa* Westwood.
Calosoter *amenetus* Walker.

- Anastatus* *cyaneicollis* Ashmead; M. O.
stantoni Ashmead; M. O.
Eupelmus *albipes* Ashmead; M. O.

Subfamily II. ENCYRTINÆ.

- Taftia* *prodeniae* Ashmead; M. O.
Howardiella *tarsata* Ashmead; M. O.
Ooencyrtus *papilionis* Ashmead; M. O.
Coccidenencyrtus *manilae* Ashmead; M. O.
Aphidencyrtus *pallidipes* Ashmead;
 M. O.
Aphytus *albilavatus* Ashmead; M. O.
Apterencyrtus *pulchricornis* Ashmead;
 M. O.
Exoristobia *philippinensis* Ashmead;
 M. O.

Family LXIX. PTEROMALIDÆ.

Subfamily IV. SPALANGIINÆ.

- Nesospalangia* *aptera* Ashmead; M. O.

Family LXX. ELASMIDÆ.

- Elasmus* *philippinensis* Ashmead; M. O.

Family LXXI. EULOPHIDÆ.

- Closterocerus* *brownii* Ashmead; M. O.
Asecodes *elasmi* Ashmead; M. O.

Subfamily II. APHELININÆ.

- Aspidiotiphagus* *aleyrodis* Ashmead;
 M. O.

Subfamily III. TETRASTICHINÆ.

- Tetrastichus* *flavobasalis* Ashmead; M. O.
philippinensis Ashmead;
 M. O.
Tetrastichoides *brownii* Ashmead; M. O.
manilensis Ashmead;
 M. O.

- Ceranisus* *incertus* Ashmead; M. O.
Gyrolasia *tagala* Ashmead; M. O.
Hyperteles *flavipes* Ashmead; M. O.

Subfamily IV. ELACHERTINÆ.

- Nesolynx* *flavipes* Ashmead; M. O.
Euplectrus *manilae* Ashmead; M. O.
philippinensis Ashmead;
 M. O.

Superfamily VIII. ICHNEUMONOIDEA.

Family LXXIV. EVANIIDÆ.

Subfamily I. EVANIDINÆ.

Evania annulipes Ashmead; M. O.
appendigaster Linnaeus.
impressa Schletterer.
verrucosa Schletterer.

Family LXXVI. ICHNEUMONIDÆ.

Subfamily I. ICHNEUMONINÆ.

Tribe II. *Ichneumonini*.

Cratichneumon manilae Ashmead; M. O.
Elasmognatha cephalotes Ashmead;
M. O.

Subfamily II. CRYPTINÆ.

Tribe II. *Phygadeuonini*.

Apsilops nigriceps Ashmead; M. O.
Isotima albicincta Ashmead; M. O.
albifrons Ashmead; M. O.
cincticornis Ashmead; M. O.

Tribe III. *Hemitelini*.

Chriodes oculatus Ashmead; M. O.
Strepsimallus bicinctus Ashmead; M. O.
Paraphylax fasciatipennis Ashmead;
M. O.
flavus Ashmead; M. O.
Acanthohemiteles benjamini Ashmead;
M. O.
Astomaspis metathoracica Ashmead;
M. O.
Microeryptus praepes Bingham,
Otaeustes alboannulus Ashmead; M. O.
Bathythrix striatus Ashmead; M. O.
Diatora prodeniae Ashmead; M. O.

Tribe VI. *Cryptini*.

Agrothereutes albicoxis Ashmead; M. O.
albipalpis Ashmead; M. O.
nigritarsis Ashmead;
M. O.
unifasciatus Ashmead;
M. O.

Agrothereutes verticalis Ashmead; M. O.
Chromocryptus albomaculatus Ashmead; M. O.

Tribe VII. *Mesostenini*.

Mesostenus leucozonatus Ashmead;
M. O.
Mesostenoideus literatus Brullé.
octozonatus Ashmead;
M. O.

Subfamily III. PIMPLINÆ.

Tribe III. *Lissonotini*.

Atropha clypearia Ashmead; M. O.
Amauromorpha metathoracica Ashmead; M. O.

Tribe IV. *Pimplini*.

Xanthopimpla kriegeri Ashmead; M. O.
Pimpla punctum Brullé.
Colpomeria flava Ashmead; M. O.

Tribe V. *Xoridini*.

Kriegeria heptazonata Ashmead; M. O.

Subfamily IV. TRYPHONINÆ.

Tribe X. *Metopini*.

Metopius browni Ashmead; M. O.

Subfamily V. OPHIONINÆ.

Tribe II. *Ophionini*.

Enicospilus ashbyi Ashmead; Ashby.
pacificus Ashmead; M. O.

Tribe IV. *Anomalini*.

Atrometus minutus Ashmead; M. O.

Tribe IVa. *Nesomesochorini*.

Nesomesochorus oculatus Ashmead;
M. O.

Tribe V. *Campoplegini*.

Charops longiventris Ashmead; M. O.
papilionis Ashmead; M. O.

Tribe VIII. *Mesochorini.*

Mesochorus philippensis Ashmead; M. O.

Tribe IX. *Porizonini.*

Leptopygus stangli Ashmead; P. L. Stangl.

Temelucha philippensis Ashmead; M. O.

Tribe X. *Pristomerini.*

Pristomerus flavus Ashmead; M. O.

Family LXXVII. ALYSIIDÆ.

Subfamily II. ALYSIINÆ.

Tribe I. *Alysini.*

Aclisis pleuralis Ashmead; M. O.

Goniarcha malayensis Ashmead; M. O.

Tribe II. *Alloenini.*

Asobara försteri Ashmead; M. O.

Family LXXXVIII. BRACONIDÆ.

Subfamily IV. METEORINÆ.

Meteorus bacoorensis Ashmead; P. L. Stangl.

brownii Ashmead; M. O.

Subfamily V. MACROCENTRINÆ.

Tribe I. *Macrocentrini.*

Macrocentrus philippensis Ashmead; M. O.

Subfamily VI. HELCONINÆ.

Tribe I. *Helconini.*

Euscelinus manilae Ashmead; M. O.

Cenocoelius albitarsis Ashmead; M. O.

Subfamily VIII. SIGALPHINÆ.

Fornicia annulipes Ashmead; M. O.

Subfamily IX. CHELONINÆ.

Chelonus albicinctus Ashmead; M. O.
semihyalinus Ashmead; M. O.

Phanerotoma albiscapa Ashmead; M. O.
philippensis Ashmead;
M. O.

Subfamily X. AGATHIDINÆ.

Tribe I. *Agathidini.*

Cremonops collaris Ashmead; M. O.

Tribe II. *Microdini.*

Stantonia flava Ashmead; M. O.

Subfamily XI. CARDIOCHILINÆ.

Cardiochiles philippensis Ashmead;
M. O.

Subfamily XII. MICROGASTERINÆ.

Glyptapanteles manilae Ashmead; M. O.
Apanteles manilae Ashmead; M. O.

philippensis Ashmead; M. O.

Urogaster albinervis Ashmead; M. O.

opacus Ashmead; M. O.

philippensis Ashmead; M. O.

stantoni Ashmead; M. O.

Microplitis manilae Ashmead; M. O.

philippensis Ashmead;
M. O.

Pseudapanteles agilis Ashmead; M. O.

Subfamily XIV. OPIINÆ.

Eurytenes nanus Ashmead; M. O.

Opius eurytenoides Ashmead; M. O.

deceptor Ashmead; M. O.

philippensis Ashmead; M. O.

Biosteres longicaudatus Ashmead; M. O.

Subfamily BRACONINÆ.

Tribe II. *Braconini.*

Bracon alguéi Ashmead; M. O.

banksi Ashmead; Banks.

ricinicola Ashmead; M. O.

trisignatus Kirby.

vau Ashmead; M. O.

Microbracon luteipes Ashmead; M. O.	Subfamily XVII. SPATHINÆ.
Habrobracon rugosiventris Ashmead; M. O.	Tribe I. <i>Pambolini</i> .
Iphiaulax deceptor Smith. luteifrons Brullé. nigifrons Brullé.	Monolexis manilensis Ashmead; M. O.
Hemiglypus flavus Ashmead; M. O.	Tribe III. <i>Spathiini</i> .
	Spathius fuscipennis Ashmead; M. O. philippensis Ashmead; M. O.
Tribe III. <i>Euurobraconini</i> .	Family LXXIX. STEPHANIDÆ.
Brownius armatus Ashmead; M. O.	Stephanus coronator Fabricius. indicus Westwood. nigricaudatus Sichel. sulcifrons Schletterer. tarsatus Schletterer. unicolor Sichel.
Subfamily XVI. RHOGADINÆ.	Suborder II. Phytophaga Ashmead.
Tribe II. <i>Rhyssalini</i> .	Superfamily IX. SIRICOIDEA.
Rhyssalus unicolor Ashmead; M. O.	Family LXX. ORYSSIDÆ.
Tribe III. <i>Rhogadini</i> .	Oryssus maculipennis Smith.
Rhogas melanosoma Ashmead; M. O. pacificus Ashmead; M. O.	Family LXXXI. SIRICIDÆ.
Tribe IV. <i>Doryctini</i> .	Tremex nigricollis Westwood.
Ischiogonus philippensis Ashmead; M. O.	Superfamily X. SELANDRIIDÆ.
Tribe V. <i>Hecabolini</i> .	Paraselandria imitatrix Ashmead; M. O. Senoclia albocoerulea Bingham.
Hecabolus ruficeps Ashmead; M. O. rufocinctus Ashmead; M. O.	

SOME OF THE PHILIPPINE HYMENOPTERA BRED IN THE MANILA
OBSERVATORY, WITH HOSTS.

1. HYMÉNOPTERA.

PARASITES.	HOSTS.
Sycorytes philippensis Ashm.	Kradibia brownii Ashm.
Philiotypesis ficieola Ashm.	Kradibia brownii Ashm.
Closterocerus brownii Ashm.	Euplectrus manilae Ashm.
Nesolynx flavipes Ashm.	Microplitis philippensis Ashm.
Asecodes elasmi Ashm.	Elasmus philippensis Ashm.

2. DIPTERA.

Anopedioides luzonicus Ashm.	Lasioptera sp.
Exoristobia philippensis Ashm.	Exorista dispar.

3. LEPIDOPTERA.

PARASITES.	HOSTS.
<i>Chaleis albotibialis</i> Ashm.	<i>Erionota thrax</i> Linn.
<i>Chaleis banksi</i> Ashm.	<i>Ticera castanea</i> Sevinhoe.
<i>Chalcis prodeniae</i> Ashm.	<i>Prodenia</i> sp.
<i>Anastatus stantoni</i> Ashm.	<i>Padraona chrisozona</i> Plotz.
<i>Eurytoma manilae</i> Ashm.	Eggs of <i>Attacus atlas</i> Linn.
<i>Taftia prodeniae</i> Ashm.	<i>Prodenia littoralis</i> Boisd.
<i>Ooencyrtus papilionis</i> Ashm.	Eggs of <i>Papilio alphenor</i> Cram.
<i>Coccidencyrtus manilae</i> Ashm.	<i>Plusia eriosoma</i> Doubl.
<i>Exoristobia philippinensis</i> Ashm.	<i>Orgyia postica</i> Walk.
<i>Euplectrus manilae</i> Ashm.	<i>Papilio alphenor</i> Cram.
<i>Euplectrus philippinensis</i> Ashm.	<i>Eucleid</i> sp.
<i>Charops longiventris</i> Ashm.	<i>Papilio rumanzovia</i> Esch.
<i>Charops papilionis</i> Ashm.	<i>Papilio agamemnon</i> Linn.
<i>Meteorus brownii</i> Ashm.	<i>Noctuid</i> sp.
<i>Apanteles philippinensis</i> Ashm.	<i>Plusia eriosoma</i> Doubl.
<i>Apanteles stantoni</i> Ashm.	<i>Pieris ida</i> Cram.
<i>Urogaster philippinensis</i> Ashm.	<i>Parasa lorquinii</i> Reak.
<i>Urogaster stantoni</i> Ashm.	<i>Tortricid</i> sp.
<i>Microplitis manilae</i> Ashm.	<i>Prodenia littoralis</i> Boisd.
<i>Microplitis philippinensis</i> Ashm.	Melopsilus acteus Cram. <i>Chaerocampa oldenlandiae</i> Fab.
<i>Rhyssalus unicolor</i> Ashm.	<i>Tineid</i> sp.

4. HEMIPTERA.

<i>Hadronotus philippinensis</i> Ashm.	Eggs of <i>Leptoglossus membranaceus</i> Fabr.
<i>Telenomus catacantha</i> Ashm.	<i>Catacanthus cerrenoi</i> Le Guillon.
<i>Aspidiotiphagus aleyrodis</i> Ashm.	<i>Aleyrodes</i> sp.